

METALOSATE® COPPER

Amino Acid Chelate Soluble Powder Foliar Fertilizer

Approved for
ORGANIC
Production

GUARANTEED ANALYSIS:

Total Nitrogen (N)	3.5%
3.5% water soluble organic nitrogen	
Copper (Cu).....	16.0%
Derived from Copper Amino Acid Chelate	

GENERAL INFORMATION:

Amino acid chelated **Metalosate® Copper** soluble powder is designed for foliar application on plants to prevent or correct nutrient deficiencies that may limit crop growth and yields. It is water soluble and nontoxic to plants when applied as directed. For best results, apply **Metalosate® Copper** soluble powder according to recommendations based on plant tissue or soil analysis.

DIRECTIONS:

Apply by a spray method and in an adequate amount of water that will provide complete coverage of the plants. Use of a non-ionic wetting agent may improve spray coverage of certain hard to wet plants.

Metalosate® Copper soluble powder can be included in a regular spray program on crops. Consult with an **Albion®** representative on compatibility with other spray materials.

The rate of application will depend on the crop, stage of growth, and severity of deficiency. The maximum recommended rates are for mature, full-sized plants. Reduce the rates proportionately when spraying smaller plants.

Metalosate® Copper soluble powder may be used on legumes, grain crops, root crops, cucurbits, cole crops, leafy vegetables, woody and herbaceous ornamentals, deciduous fruits, vine crops, tropical and subtropical fruits, and many other crops.

Albion® and Metalosate® are trademarks Albion Laboratories, Inc. Patent information: www.albionplantnutrition.com/patents

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfc.org/metals.htm>

Manufactured by: **Albion Laboratories, Inc.**
67 South Main Street
Layton, Utah 84041 U.S.A.

Distributed by: **Chemseed Pty Ltd**
Units 7-8, Warehouse J
Flemington Markets NSW 2129
Phone: 0419.684.117

RECOMMENDATIONS:

Field Crops and Vegetables: Apply 0.15 to 0.3 kg per hectare during the periods of rapid growth or nutritional stress. The application may be repeated 2 or more times through the growing season.

Tree Crops: Make an application of 0.15 to 0.3 kg per hectare after the beginning of active growth. The application may be repeated at 2 to 4 week intervals through the growing season.

Grapes and Berries: Make an application of 0.15 to 0.3 kg per hectare after active growth begins. The application may be repeated at intervals of one week or more through the vegetative-growth period.

LIMITED WARRANTY:

Seller's and Manufacturer's obligation is limited to replacement of defective product. **NEITHER SELLER NOR MANUFACTURER SHALL BE LIABLE FOR ANY INJURY, LOSS, OR DAMAGE DIRECTLY OR CONSEQUENTLY ARISING OUT OF THE MISUSE OR INABILITY TO USE THE PRODUCT. ALL OTHER WARRANTIES EXCEPT IN WRITING FROM THE MANUFACTURER, WHETHER EXPRESSED OR IMPLIED ARE HEREBY DISCLAIMED.**

Reseal package between mixings to maintain freshness.

This product has been
verified to comply with 7 CFR, Part 205
United States Department of Agriculture
National Organic Program

WARNING: Harmful if swallowed, causes serious eye irritation, Very toxic to aquatic life with long lasting effects. Do not eat, drink or smoke when using this product. Wear protective gloves/protective clothing/eye protection/face protection. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention. Collect spillage. IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell. Rinse mouth. Dispose of contents/container in an approved waste disposal plant.

**READ LABEL BEFORE USE.
KEEP OUT OF REACH OF CHILDREN.**

**IF MEDICAL ADVICE IS NEEDED:
CONTACT THE NATIONAL POISONS CENTRE,
PHONE NZ 0800 POISON (0800 764 766)
OR AUSTRALIA 13 11 26
OR A DOCTOR. HAVE PRODUCT CONTAINER
OR LABEL ON HAND.**



label version
AU05/10/2019

